



Public Health  
Agency of Canada

Agence de la santé  
publique du Canada



**National Integrated Enteric Disease Surveillance Program**  
Sample Collection, Preparation & Laboratory Methodologies  
January 2010

# Table of Contents

---

---

Table of Contents.....	i
1. Sample Collection.....	1
Retail Meat.....	1
Agriculture.....	1
Untreated Surface Water.....	2
Human.....	2
2. Sample Preparation.....	3
Retail Meat.....	3
Agriculture.....	3
Untreated Surface Water.....	4
3. Laboratory Methodologies.....	5
3.1 Campylobacter.....	5
Retail Meat.....	5
Agriculture.....	5
Untreated Surface Water.....	5
Subtyping (Retail, Agriculture, Water).....	6
Human.....	7
Subtyping (Human).....	8
3.2 Salmonella.....	8
Retail Meat.....	8
Agriculture.....	9
Untreated Surface Water.....	9
Subtyping (Retail, Agriculture, Water).....	9
Human.....	11
Subtyping (Human).....	12
3.3 Pathogenic E. coli.....	14
Retail Meat.....	14
Agriculture.....	14
Untreated Surface Water.....	15
Subtyping (Retail, Agriculture).....	15
Human.....	16
Subtyping (Human).....	17
3.4 Yersinia.....	18
Retail Meat.....	18
Agriculture.....	18
Untreated Surface Water.....	20
Subtyping (Retail, Agriculture, Water).....	19
Human.....	20
Subtyping (Human).....	21
3.5 Listeria.....	22
Retail Meat.....	22
Agriculture.....	22
Subtyping (Retail, Agriculture).....	22

3.6	Giardia and Cryptosporidium .....	23
	Agriculture .....	23
	Untreated Surface Water .....	23
	Human .....	23
3.7	Viruses .....	25
	Agriculture .....	25
3.8	Sources Cited .....	26
	Appendix .....	28
	C-EnterNet Retail Sample Collection Overview .....	28
	C-EnterNet Agriculture Sample Collection Overview .....	29
	C-EnterNet Surface Water Collection Overview .....	30
	C-EnterNet Human Sample Collection Overview .....	31

# 1. Sample Collection

---

## Retail Meat

C-EnterNet's retail surveillance component involves a random and systematic approach to derive estimates of the pathogen levels on raw meat available to the consumer at the grocery store level. The sampling framework is based on a census of the retail grocery store outlets operating within the sentinel site. This includes large and medium sized chain stores as well as independently owned butcher and market shops. Each week, three chain stores and one independent outlet are randomly selected from the store census. A C-EnterNet field sampler anonymously visits the selected stores and purchases one ground beef, one chicken breast and one pork chop package. The meat package purchased is required to be fresh (not previously frozen) and at least 500-600 g to ensure sufficient amount of sample for bacterial testing. Twelve meat samples per week are collected with a total of 600 samples per year (50 weeks/year sampling). Samples are shipped at refrigeration temperature (4-6°C) to a laboratory for primary isolation, enumeration and subtyping.

## Agriculture

C-EnterNet's agriculture surveillance component involves the sampling of food-animal operations (swine, beef, dairy and poultry) within the sentinel site. Farm selection and visitation is performed through a contracted third party to ensure producer confidentiality. In general, 30 farms for each commodity are randomly selected for a single visitation during the calendar year. Each month, approximately 2 farms per commodity are visited. At each farm visit, a short farm management questionnaire is administered and manure samples are collected.

Three pooled fresh fecal samples representing different age or production groups and a pooled stored manure sample are collected at each visit. In total, 120 samples are collected per year for each commodity. Samples are shipped in coolers at refrigeration temperatures (4-6°C) for primary isolation and subsequent subtyping.

### **Fresh Fecal Sampling Protocol**

Fresh fecal samples from five individual animals are collected and put into one large plastic bag and gently mixed. A sterile scoop is used to transfer the pooled sample to specimen containers.

### **Stored Manure Sampling Protocol**

#### a) Liquid Pit Sampling:

A sampling pole and bottle holder are used to collect three sub-samples from three locations around the pit, and if possible, up to two depths (i.e. the top 1/3, and mid depth of the storage), for a total of 6 sub-samples. The sub-samples are poured into a sampling bucket with a clean plastic liner and mixed to create the pooled sample.

b) Dry Pile Sampling:

A sterile sampling scoop is used to select sub-samples from five different locations in the stored manure area. These sub-samples are then added to a sampling bucket with a clean plastic liner bag and mixed to create a pooled sample.

## **Untreated Surface Water**

C-EnterNet's water sampling and analysis framework includes a systematic sampling frame of the Grand River watershed in the sentinel site in the Region of Waterloo, Ontario. The Grand River watershed is sampled twice per month at five locations upstream of the regional drinking water intake. The approximate number of samples per year is 120. At each sample collection, field staff collect water samples using 1 L sterile sampling bottles - containing sodium thiosulfate - and an extendable sampling pole to collect from a fast flowing portion of the river.

Field staff deliver coolers with samples to Laboratory Services Division, University of Guelph where bacteriology testing is conducted. Duplicate samples are sent to Hyperion Ltd. for parasitology (*Cryptosporidium* and *Giardia*) testing (C-EnterNet Watershed Data Collection Overview-Appendix; Ministry of Environment Water Treatment Laboratory Sample Collection Protocol).

## **Human**

C-EnterNet's human stool samples are collected through the existing passive surveillance system in Ontario. A passive surveillance system is based on case presentation to physicians in which a stool specimen is requested and then submitted to a private hospital or public health laboratory for testing for enteric pathogens that are reportable in the province. A physician can request three types of stool samples for analysis (i.e. parasite and ova, culture and sensitivity, or viral). For the case to be captured by the public health reporting system, either the laboratory or the physician must report the positive isolation directly to a local health authority, who will then report the case to the province, and subsequently that case is reported to the national level.

C-EnterNet works with the provincial public health laboratory system, three private laboratories and the regional hospital laboratory that serve Sentinel Site 1, to ensure that the positive isolations of reportable enteric pathogens that have been initially identified are forwarded to the Ontario Agency of Health Protection and Promotion's Public Health Laboratory in Toronto and the Public Health Agency of Canada's National Microbiology Laboratory in Winnipeg for subtyping.

An enhanced standardized questionnaire for endemic enteric disease cases was among the first new epidemiological tools to result from C-EnterNet's partnership with the local public health department at its pilot sentinel site in the Region of Waterloo, Ontario. The questionnaire, originally based on the one used at the CDC FoodNet site at the Minnesota Department of Health, is designed to secure additional risk factor information for each enteric case, to support further epidemiological investigation and C-EnterNet's surveillance. The questionnaire data are manually entered into iPHIS and EpiData for further analysis.

## **2. Sample Preparation**

---

### **Retail Meat**

1. Meat packages are removed from the cooler and placed on a clean surface. The external meat package surface is wiped with sterile gauze and 70% ethanol. The meat package is then aseptically opened using sterilized scissors. The external surface of the wrapping is prevented from touching the meat samples to avoid possible contamination of meat from outside sources.
2. Sterile tongs are used to remove the meat samples from its package. Several pieces of meat are cut from different areas of the meat sample to make a representative sample weighing 50 g. This is placed into a prepared stomacher bag with selective enrichment broth specific to each of the following primary detection methods:
  - a. 50 g of meat sample into 450 ml of Buffered Peptone Water (BPW) used for primary isolation of:
    - VTEC
    - *E. coli* Coliform counts
    - *Salmonella* spp.
  - b. 50 g of meat sample into 250 ml of Bolton Broth used for primary isolation of :
    - *Campylobacter* spp.
  - c. 50 g of meat sample into 450 ml of *Listeria* Enrichment Broth (LEB) for primary isolation of:
    - *Listeria monocytogenes*
  - d. 50 g of PORK meat sample into Luria Bertani-Bile Salt Irganon (LB-BSI)
    - *Yersinia enterocolitica*
3. Samples are then massaged or stomached in a stomacher bag for 2 minutes.

### **Agriculture**

1. Sample containers are placed on a clean surface, and then weighed.
2. Sample containers are then carefully opened to avoid spillage due to gas build-up during transport.
3. Sterile tongue depressors are used to mix samples gently and to transfer the samples into prepared stomacher bags with selective enrichment broth specific to each of the following primary detection methods:

## Swine samples

- a. 25 g of fecal sample into 100 ml of Buffered Peptone Water (BPW).  
5 ml of the BPW mix for primary isolation of:
  - *Yersinia enterocolitica*
- b. More BPW is added to the remaining BPW mix to obtain a 1:10 dilution which is used for primary isolation of
  - *Salmonella* spp.
  - *E. coli* 0157:H7
  - *Campylobacter* spp.
  - Original fecal sample preservation
- c. 1 ml of BPW mix into 9 ml (1:10) of Bolton broth for primary isolation of:
  - *Campylobacter* spp.

## Bovine and poultry samples

- a. 10 g of fecal sample into 90 ml (1:10) of Buffered Peptone Water (BPW) used for primary isolation of:
  - *Salmonella* spp.
  - *E. coli* 0157:H7
  - *Campylobacter* spp.
  - Original fecal sample preservation
- b. 1 ml of BPW mix into 9 ml (1:10) of Bolton broth for primary isolation of:
  - *Campylobacter* spp.

4. Massage or stomach samples in a stomacher for 30 sec.

## Untreated Surface Water

1. Within 48h of collection, start processing samples. Remove bottles from refrigerator (4<sup>0</sup> C) on a clean surface.
2. Using proper aseptic techniques remove appropriate amount of water, filter through a 0.45µm filter (Millipore filters) and place filter membrane into selective enrichment specific to each of the following primary detection methods:
  - a. 1000ml for *Yersinia enterocolitica*
  - b. 1000ml water for *Salmonella* spp.
  - c. 1000ml water for *Campylobacter* spp.
  - d. 1000ml water for *E. coli* 0157
  - e. 1000ml water for *E. coli* 0157 enumeration

## **3. Laboratory Methodologies**

---

### **3.1 *Campylobacter***

#### **Retail Meat**

##### **Culture Detection**

Samples are processed and enriched in Bolton Broth at 42°C. Following 48 hrs incubation at 42°C under microaerophilic conditions, samples are streaked onto selective agar plates. Suspect colonies are identified using microscopy. Typical *Campylobacter* colonies are confirmed and speciated using biochemical methods.

#### **Agriculture**

##### **Culture Detection**

Method # 1 (implemented in 2008)

Bovine and poultry fecal samples are processed by transferring 10 g of feces in 90 ml of Buffered Peptone Water (BPW). For porcine fecal samples 25 g of feces are transferred in 225 ml of BPW. Then 1 ml of the BPW mix is transferred in 9 ml of Hunt's Enrichment Broth (HEB) and incubated in a microaerophilic atmosphere at 35°C for 4 hrs. After this first incubation, 36 µL of sterile cefoperazone is added to the HEB. Tubes are then incubated in microaerophilic condition at 42°C for 20 to 24hrs. A loop of the incubated HEB is then plated onto modified cefoperazone charcoal deoxylate agar (mCCDA) agar plate. Plates are incubated at 42°C in microaerophilic conditions for 72 hrs. Suspect colonies are further tested using a second mCCDA plate, gram stain, dark field microscopy, and Mueller Hinton Blood (MHB) agar. Typical *Campylobacter* colonies are confirmed using catalase and oxidase tests. Hippurate hydrolysis and indoxyl acetate hydrolysis tests and cephalothin susceptibility tests are performed to speciate suspected *Campylobacter* into *C. jejuni*, *C. coli* and *C. lari*.

Method #2 (utilized in 2006-2007)

Samples are processed by transferring 1g of manure into 9 ml of Bolton broth. Samples are incubated at 42°C for 48 hrs in microaerophilic conditions, and plated onto modified cefoperazone charcoal deoxylate agar (mCCDA) agar plate. Plates are incubated at 42°C in microaerophilic conditions for 24 to 48 hrs. Suspect colonies are further tested using a second mCCDA plate, gram stain, dark field microscopy, and Mueller Hinton Blood (MHB) agar. Typical *Campylobacter* colonies are confirmed as in method #1.

#### **Untreated Surface Water**

##### **Culture Detection**

A 1L water sample is filtered using 0.45µm filter membrane. Filter samples are enriched in Bolton broth, incubated at 42°C for 48h. After the first enrichment incubation period, samples are plated onto modified cefoperazone charcoal deoxycholate agar (mCCDA) agar and incubated in microaerophilic conditions at 42°C for 3 days. MCCDA Plates are examined after 48 and 72 h. Suspect colonies are further plated on Mueller Hinton Blood



(MHB) agar and incubated under microaerophilic conditions at 37°C for 48 h. Presumptive *Campylobacter* colonies are confirmed from a fresh MHB plate using darkfield microscopy examination, catalase, oxidase, the oxygen tolerance test, and by testing for growth at 25°C. Hippurate hydrolysis and indoxyl acetate hydrolysis tests are performed to speciate suspected *Campylobacter* into *C. jejuni*, *C. coli*, or *C. lari*. Susceptibility testing is performed for presumptive *Campylobacter* spp. at LFZ-SHU and targeted subset is sent to NML for Comparative molecular testing/PFGE (LSD *Campylobacter* Isolation Method).

### **Molecular Detection**

Quantitative PCR for *Campylobacter* identification in river water samples uses primers designed by Lubeck et al (2003) together with a probe designed by the University of Waterloo (UW) NSERC Chair in Water Treatment research group. The assay is specific for the 16S rRNA gene of *C. jejuni*, *C. coli* and *C. lari*.

### **Subtyping (Retail, Agriculture, Water)**

#### **Molecular Comparative Testing and PFGE**

A subset of *Campylobacter* isolates are sent to the Public Health Agency of Canada's National Microbiology Laboratory (NML) for comparative molecular testing and PFGE testing.

### **Antimicrobial Susceptibility Testing (2005-2007)**

Antimicrobial susceptibility of *Campylobacter* is performed by LFZ AMR Laboratory, Saint-Hyacinthe Unit using a broth microdilution Sensititre™ plate (Trek Diagnostic System Ltd.). Isolates are prepared for testing by streaking colonies on Mueller Hinton agar containing 5% lysed horse blood and incubating them in a microaerophilic atmosphere for 24 hrs at 42°C. Isolated colonies are then mixed in 5 ml of Sensititre™ cation adjusted Mueller Hinton broth with TES buffer (CAMHBT) and adjust to 0.5 MacFarland. One hundred µl of this broth is then mixed with 11 ml of CAMHBT with lysed horse blood, which is then inoculated in a Sensititre™ Campy plate. After 24 hrs incubation in a microaerophilic atmosphere at 35°C plates are read visually using Sensitouch™ giving minimum inhibitory concentrations (MIC) values (since 2009, Sensitouch has been replaced by Vizion which also permits visual reading using MIC values). The following are the antimicrobials tested: azithromycin, ciprofloxacin, clindamycin, erythromycin, florfenicol, gentamicin, nalidixic acid, telithromycin and tetracycline. The Canadian Integrated Program for Antimicrobial Resistance Surveillance (CIPARS) breakpoints are used to determine resistance profiles and can be found at : [http://www.phac-aspc.gc.ca/ciparspicra/analys\\_e.html](http://www.phac-aspc.gc.ca/ciparspicra/analys_e.html).

### **Enumeration - Most Probable Number (MPN)**

The Most Probable Number (MPN) method is used for retail samples that are positive for *Campylobacter*. This method estimates the number of *Campylobacter* species per millilitre or gram of sample using three tube MPN dilutions. Fifty grams of meat are stored under microaerophilic conditions at 4 °C for the MPN test. Positive samples are then retrieved, and added to Hunts Enrichment Broth (HEB); processed and serial dilutions are prepared. Tubes are incubated at 36°C for 4 hrs and following incubation cefaperazone is added to each sample tube and incubated at 42°C for 24 hrs under microaerophilic conditions. Then, each sample is streaked onto selective agar plates. Suspect colonies are identified using microscopy. Typical *Campylobacter* colonies are

confirmed using biochemical methods. MPN is calculated using the table from the FDA Bacteriological Analytical Manual (<http://www.cfsan.fda.gov/~ebam/bam-toc.html>).

## **Human**

### **LAB A**

Specimens are collected and mixed with Enteric Plus transport media before arriving at the laboratory. Once at the laboratory, a sterile swab is used to obtain a homogenized sample, which is streaked for isolated colonies onto Karmali blood-free agar. Plates are incubated at 42°C for 48 hrs in microaerophilic conditions using a CO<sub>2</sub>-N<sub>2</sub>-O<sub>2</sub> percentage ratio of 10:80:10. Grey suspect colonies are sub-plated onto sheep blood agar for speciation testing. In some cases the original Karmali plate is used for biochemical testing. Isolates are subject to hippurate, catalase, TSI for H<sub>2</sub>S testing and cephalothin and nalidixic acid susceptibility testing.

### **LAB B**

Specimens are collected and mixed with Cary-Blair transport media before arriving at the laboratory. Once at the laboratory, a sterile swab is used to obtain a homogenized sample which is streaked for isolated colonies onto *Campylobacter* blood-free agar. Plates are incubated at 42°C for 48 hrs in microaerophilic conditions using a CO<sub>2</sub>-N<sub>2</sub>-O<sub>2</sub> percentage ratio of 10:85:5. Suspect colonies are subcultured onto *Campylobacter* blood-free agar or sheep blood agar for speciation testing. In some cases the original *Campylobacter* blood-free plate is used for biochemical testing. Isolates are subject to hippurate, oxidase, and gram stain testing.

### **LAB C**

Specimens are collected and mixed with Cary-Blair transport media before arriving at the laboratory. Once at the laboratory, a sterile swab is used to obtain a homogenized sample which is streaked for isolated colonies onto *Campylobacter* blood-free agar. Plates are incubated at 42°C for 48 to 72 hrs in microaerophilic conditions. Suspect colonies are subcultured onto a blood agar plate for speciation testing. In some cases the original *Campylobacter* blood-free plate is used for biochemical testing. Isolates are subject to hippurate, oxidase, and gram stain testing.

### **LAB D**

Specimens are collected in a container with Cary-Blair transport media and mixed by the patient before arriving at the laboratory. Once at the laboratory, a sterile swab is used to obtain a homogenized sample, which is streaked for isolated colonies onto *Campylobacter* blood-free agar. Plates are incubated at 42°C for 36 to 60 hrs in microaerophilic conditions using Oxoid Campygen pack®. Suspect colonies are subcultured onto a blood agar plate for speciation testing. Isolates are tested by hippurate, oxidase, and gram stain.

## **Ontario Agency for Health Protection and Promotion, Public Health Laboratory – London and Toronto locations**

Stool specimens are collected and mixed with Cary-Blair transport medium at the time of collection. Once received in the laboratory, a sterile swab is used to obtain a representative sample, which is streaked onto a charcoal-based selective medium (CSM) for *Campylobacter* isolation. Plates are incubated at 42°C for 18 - 48 hours in

microaerophilic conditions using a CO<sub>2</sub>-N<sub>2</sub>-O<sub>2</sub> percentage ratio of 10:85:5. A liquid enrichment broth (LEM) is also inoculated upon receipt of the specimen and incubated at the same temperature as the plate medium for 18-24 hours. A loopful of the enrichment solution is streaked for isolated colonies onto a CSM agar plate and incubated at 42°C for 18-48 hours. Suspect colonies are sub-cultured onto three Columbia Blood Agar plates and incubated for 18-48 hours at different temperatures for speciation testing. The first Columbia Blood Agar plate is incubated at 25°C (aerophilic), the second plate at 35°C and the third plate at 42°C (microaerophilic). Isolates are subject to Hippurate hydrolysis, Oxidase production test, Gram stain reaction, Catalase test, Indoxyl acetate hydrolysis test and Nalidixic acid/Cephalothin susceptibility testing. Additional tests to aid in identification may include: Nitrate Reduction Test, Urea hydrolysis, H<sub>2</sub>S (TSI) production and tolerance to 1% glycine.

### **Subtyping (Human)**

All *Campylobacter* species isolates from the Region of Waterloo including *C. jejuni* and *C. coli* are tagged as a C-EnterNet sample and sent to the Public Health Laboratory - Toronto for further testing. Isolates are also forwarded to the National Microbiology Laboratory in Winnipeg, Manitoba for experimental comparative molecular subtyping.

### **Antimicrobial susceptibility Testing**

Antimicrobial susceptibility testing of *Campylobacter* was not performed in 2008.

## **3.2 Salmonella**

### **Retail Meat**

#### **Culture Detection**

Retail meat samples are screened by a modification of MFLP-75 primary isolation method of the *Compendium of Analytical Methods, Health Protection Branch, and Methods of Microbiological Analysis of Food, Government of Canada* is utilized. In order to minimize possible false negative results, a secondary enrichment followed by Modified Semi-solid Rappaport Vassiliadis (MSRV) plating. This method is applicable to viable, motile *Salmonella* species found in meat products.

Samples are pre-enriched in a non-selective broth (buffered peptone water) at 35°C. One millilitre is transferred to 9 ml of tetrathionate brilliant green (TBG) broth. An MSRV plate is inoculated with TBG broth and incubated at 42°C for 72 hrs. The plates are examined at 48 and 72 hours and suspect colonies are further screened and confirmed using biochemical methods.

## **Agriculture**

### **Culture Detection**

A modification of MFLP-75 primary isolation method of the *Compendium of Analytical Methods, Health Protection Branch, Methods of Microbiological Analysis of Food, Government of Canada* is used for the detection of *Salmonella* in manure. *Salmonella* spp. can be detected in Modified Semi-solid Rappaport Vassiliadis (MSRV) media at a temperature of 42°C.

Samples are pre-enriched in a non-selective broth (buffered peptone water) at 35°C. In parallel, 0.1ml of the pre-enrichment broth is inoculated into MSRV. Samples are incubated at 42°C for 24 to 72 hrs and examined at 24 hr intervals. Suspect colonies are further screened on MacConkey, Triple Sugar Iron (TSI) and Urea agar plates. Further testing of presumptive colonies is performed with the Kovac's Indole test and the slide agglutination test using the *Salmonella* O antiserum Poly A-I and V1.

## **Untreated Surface Water**

### **Culture Detection**

A modification of *Salmonella* isolation protocol of the USDA Food Safety and Inspection Service (FSIS) method for isolation and detection of *Salmonella* (MLG 4.04) is used. Suspect salmonellas are confirmed with the Health Canada method for the isolation and identification of *Salmonella* MFHPB-20.

A one litre water sample is filtered using 0.45µm filter membrane. Filtered samples are pre-enriched in a non-selective broth (buffered peptone water) at 35±2°C for 20-24h. Following pre-enrichment, samples are further enriched with selective enrichment broths (Rappaport Vassiliadis and tetrathionate) and incubated at 42±0.5°C for 22-24h. From each selective enrichment broth, samples are transferred to Xylose Lysine Tergitol 4 (XLT4) and Brilliant Green sulfa (BGS) agar. Suspect colonies are further screened on Triple Sugar Iron (TSI), Lysine Iron Agar (LIA), Christensen's Urea Agar, and streaked onto MacConkey Agar for further incubation at 35±2°C for 18-24h. Confirmation is performed by serology with polyvalent antiserum and API 20E test strips. Presumptive *Salmonella* isolates are sent to the LFZ-Guelph where the *Salmonella* Typing Laboratory (STL) performs serotyping and phage typing and the Canadian Integrated Program for Antimicrobial Resistance Surveillance (CIPARS) Guelph laboratory performs susceptibility testing (LSD-MID 235 Rev. 2 and MID 1021 isolation and identification of *Salmonella* from meat, poultry and egg products USDA MLG 4.04).

## **Subtyping (Retail, Agriculture, Water)**

### **Serotyping**

*Salmonella* isolates are transferred on agar slants to the Public Health Agency of Canada - Laboratory for Foodborne Zoonoses (LFZ), Office International des Epizooties Reference Laboratory for Salmonellosis, *Salmonella* Typing Laboratory Guelph, Ontario for serotyping and phagetyping. Briefly, the serotyping procedure involves determining the O or somatic antigens of the *Salmonella* isolates using slide agglutination Ewing (1986). The H or flagellar antigens are identified using a microtechnique Shipp and Rowe (1980) that employs microtitre plates. The antigenic formulae of Le Minor and Popoff (2001) are used to identify and name the serovars.

## Phagetyping

Phage typing is performed for *Salmonella* Enteritidis, S. Hadar, S. Heidelberg and S. Typhimurium. The standard phagetyping technique described by Anderson and Williams (1956) is followed. *Salmonella* Enteritidis isolates are phagetyped with typing phages obtained from the International Centre for Enteric Phage Typing (ICEPT), Central Public Health Laboratories, Colindale, UK Ward (1987) via the Enteric Diseases Program, National Microbiology Laboratory (NML), Public Health Agency of Canada, Winnipeg, Manitoba. *Salmonella* isolates are categorized according to reaction with phage. Isolates not fitting into a recognized phage type are labelled atypical and isolates that do not react with any of the phages provided are considered untypable (Laboratory for Foodborne Zoonoses *Salmonella* Serotyping and Phage typing Manual, SLP-001, Version 4). The phage typing scheme and phages for *Salmonella* Typhimurium developed by Callow (1959) and further extended by Anderson (1964) and Anderson et al. (1977) were obtained from the ICEPT via the NML. The *Salmonella* Heidelberg phagetyping scheme and phages were supplied by the NML (Demczuk et al, 2003).

## PFGE Testing

PFGE testing is performed at the Laboratory for Foodborne Zoonoses, Guelph (*E. coli* Lab) according to standardized protocols developed by PulseNet USA and in use by PulseNet Canada. For *Salmonella*, two restriction enzymes, *Xba*I and *Bln*I, are used to obtain two separate patterns for each strain. The resulting tiff files are uploaded to the national database at the NML to obtain the national designation for the two separate patterns.

## Antimicrobial Susceptibility Testing (2005-2007)

Antimicrobial susceptibility testing is performed by the CIPARS Guelph Laboratory at the Laboratory for Foodborne Zoonoses using the Sensititre<sup>®</sup> Automated Microbiology System (Trek Diagnostic Systems Ltd.). *Salmonella* isolates are prepared by streaking the cultures onto Mueller Hinton Agar for isolated colonies, and then for growth, and incubated for 18 to 24 hrs at 35°C. Each culture is prepared in Sensititre<sup>®</sup> demineralized water to a 0.5 McFarland standard and then inoculated into Sensititre<sup>®</sup> Mueller Hinton Broth with TES. The NARMS Gram negative plate with the CMV1AGNF panel is inoculated with the broth suspension and is incubated for 18 hrs at 37°C. Each plate is then read by the Sensititre<sup>®</sup> AutoReader to determine the minimum inhibitory concentration. The following are the antimicrobials tested: amikacin, ampicillin, amoxicillin/clavulanic acid, cefoxitin, ceftiofur, ceftriaxone, chloramphenicol, ciprofloxacin, gentamicin, kanamycin, nalidixic acid, streptomycin, sulfisoxazole, tetracycline, trimethoprim/sulfamethoxazole. CIPARS breakpoints are used to determine resistance profiles and can be found at:

[http://www.phac-aspc.gc.ca/ciparspicra/analys\\_e.html](http://www.phac-aspc.gc.ca/ciparspicra/analys_e.html).

## Enumeration - Most Probable Number (MPN)

The Most Probable Number method is only used for retail samples positive for *Salmonella* from the detection method. This method estimates the number of *Salmonella* species per ml or g of sample using three tube MPN dilutions. Fifty millilitres of the original processed sample is stored for MPN testing to prepare serial dilutions. Tubes are incubated at 35°C for 24 hrs. After incubation, 1 ml of each tube is transferred to TBG broth, and 0.2ml of potassium iodine is added to each new tube and incubated at 42°C for 24 hrs. Two different selective media plates are streaked and incubated at

35°C for 24 hrs. Suspect colonies are confirmed with biochemical methods. The MPN is calculated using the MPN table from the FDA Bacteriological Analytical Manual (<http://www.cfsan.fda.gov/~ebam/bam-toc.html>).

## **Human**

### **LAB A**

Specimens are collected and mixed with Enteric Plus transport media before arriving at the laboratory. Once at the laboratory, the Enteric Culture Manual method for *Salmonella* spp. is used. A sterile swab is used to obtain a homogenized sample, which is streaked for isolated colonies onto MacConkey agar and *Salmonella-Shigella* (SS) agar. Samples are incubated at 35°C for 18 to 24 hrs. Once at the laboratory, a portion of the homogenized sample is inoculated into Selenite F broth and incubated at 35°C for 12 hrs. The broth is subcultured onto an SS agar plate and incubated for 18 to 24 hrs. Suspect colonies are identified by non-lactose fermenting (NLF) colonies with or without H<sub>2</sub>S production on the SS plate. Sheep blood agar or MacConkey agar is used for *Salmonella* spp confirmation with triple sugar iron (TSI), OPM, antisera agglutination and Microscan biochemical panel of tests. All *Salmonella* species are sent to TRPHL for species identification.

### **LAB B**

Specimens are collected and mixed with Cary-Blair transport media before arriving at the laboratory. Once at the laboratory, a portion of the specimen solution is inoculated in Selenite F broth and incubated at 35°C for 12 hrs. At the same time, a Hektoen plate is streaked using the original specimen incubated at 35° for 18 to 24 hrs. After incubation, using a sterile swab, the enrichment solution is streaked for isolated colonies onto Hektoen agar. Samples are incubated at 35°C for 18 to 24 hrs. Non-lactose fermenting (NLF) colonies with or without H<sub>2</sub>S production on Hektoen are screened using urea, SIM and PYR. For confirmation testing, the original Hektoen plate, or a freshly subcultured sheep blood agar, is used for testing suspect isolates. Confirmation tests include Vitek 2 ID system and antisera latex agglutination test. All *Salmonella* species are sent to TRPHL for identification. Susceptibility testing is performed on all sterile sites, as per CLSI M100 using Viteck2 and Nalidixic KB. For stool samples, susceptibility testing is only performed on *S. Typhimurium* and *S. Paratyphi*. For other *Salmonella* species, susceptibility testing is done if the individual is less than 3 months of age or older than 65 year of age.

### **LAB C**

Specimens are collected and mixed with Cary-Blair transport media before arriving at the laboratory. Once at the laboratory, a portion of the specimen solution is inoculated in Selenite F broth and incubated at 36°C for 18 to 24 hrs. Using a sterile swab, the enrichment solution is streaked for isolated colonies onto Hektoen agar and MacConkey agar. Samples are incubated at 36°C for 48 hrs. Suspect colonies are identified by non-lactose fermenting (NLF) colonies with H<sub>2</sub>S production. For confirmation testing, a freshly subcultured CLED plate is used. Confirmation tests include Urea, TSI, ONPG, SIM, Phoenix ID system and an agglutination test.

### **LAB D**

Specimens are collected in a container with Cary-Blair transport media and mixed by the patient before arriving at the laboratory. Once at the laboratory, a portion of the

specimen is inoculated in Selenite F broth and incubated at 37°C for 8 to 12 hrs. At the same time, a Hektoen plate is streaked using the original specimen solution and incubated at 37°C for 18 to 24 hrs. After incubation, using a sterile swab, the enrichment solution is streaked for isolated colonies onto Hektoen agar and incubated at 37°C for 18 to 24 hrs. Non-lactose fermenting (NLF) colonies with or without H<sub>2</sub>S production are screened with TSI, SIM and Urea tests. Confirmation tests include Enterotube or the Vitek card automated ID system, PYR, Indole and an agglutination test performed from a blood agar plate. Isolates are forwarded to PHL for monotyping.

### **Ontario Agency for Health Protection and Promotion, Public Health Laboratory – London and Toronto locations**

Stool specimens are collected and mixed with Cary-Blair transport medium at the time of collection. Once received in the laboratory, a sterile swab is used to obtain a representative portion of the sample which is streaked for isolated colonies onto MacConkey Agar, Xylose-lysine-deoxycholate (XLD) Agar, and *Salmonella-Shigella* (SS) Agar. A Selenite F broth is also inoculated and all media are incubated at 36°C for 18 hours. After incubation the Selenite F enrichment broth is sub-cultured and streaked for isolated colonies onto SS Agar and incubated for the same time and environmental conditions as the primary SS Agar plate. Suspect colonies are non-lactose fermenting (NLF) colonies with or without H<sub>2</sub>S production. Typical NLF colonies are screened using TSI, ONPG-PAM with an indole strip, lysine decarboxylase and PYR test. Confirmation is performed on Blood Agar sub-cultures of screen positive isolates using API biochemical strips and slide agglutination using *Salmonella* polyvalent antiserum. Further confirmatory testing performed at the Public Health Laboratory - Toronto include using a freshly sub-cultured Blood Agar plate, MacConkey plate and motility plate (Gard plate) for testing isolates as well as a panel of biochemical tests. Final speciation of the *Salmonella* isolates involve using the freshly sub-cultured isolate to perform serotyping which consists of agglutination tests for somatic and flagellar antigens.

### **Subtyping (Human)**

As part of C-EnterNet's enhanced public health passive surveillance system, all isolates are submitted to the Public Health Laboratory - Toronto for serotyping and pulse-field gel electrophoresis (PFGE). Isolates are also forwarded to the National Microbiology Laboratory in Winnipeg, Manitoba for phagetyping where appropriate and antimicrobial susceptibility testing.

### **Serotyping**

The serotyping method is based on the following references:

- Farmer III JJ. 2003 *Enterobacteriaceae*: introduction and Identification, p. 636-653. In PR Murray, EJ Baron, MA Tenover, JH Tenover ed., *Manual of clinical microbiology*, 8<sup>th</sup> ed. American Society for Microbiology, Washington, D.C.
- Ewing WH. Edwards and Ewing's Identification of *Enterobacteriaceae*, 4<sup>th</sup> ed, pp. 181-304, 1986, Elsevier Science Publ., New York.
- Popoff MY. 2001. Antigenic Formulas of the *Salmonella* Serovars, 8<sup>th</sup> ed. WHO Collaborating Center for Reference and Research in *Salmonella*, Institut Pasteur, Paris, France.



## Phagotyping

Phagotyping is performed at the NML, Winnipeg, in the Enteric Diseases Program lab.

## PFGE Testing

The PFGE method used at the Public Health Laboratory – Toronto is based on the national standardized PulseNet Canada protocol.

One-day (24-48h) Standardized Laboratory Protocol for the Molecular Subtyping of Enteric Pathogens by Pulsed Field Gel Electrophoresis (PFGE) and Standardized Molecular Subtyping of Foodborne Bacterial Pathogens by Pulsed Field Gel Electrophoresis (Center for Disease Control and Prevention Manual –PNL 05) are followed for the PFGE testing of *Salmonella* isolates.

## Antimicrobial Susceptibility Testing

*Salmonella* isolates are forwarded to the National Laboratory for Enteric Pathogens (NLEP) for antimicrobial susceptibility testing, according to the Sensititre®: automated micro-dilution AMR test.

The following are the antimicrobials tested: amikacin, amoxicillin/clavulanic acid, ampicillin, cefoxitin, ceftiofur, ceftriaxone, cephalothin, chloramphenicol, ciprofloxacin, gentamicin, kanamycin, nalidixic acid, streptomycin, sulphamethoxazole, tetracycline, and trimethoprim/sulfamethoxazole. CIPARS breakpoints are used to determine resistance profiles and can be found at:

[http://www.phac-aspc.gc.ca/ciparspicra/analys\\_e.html](http://www.phac-aspc.gc.ca/ciparspicra/analys_e.html).

## Antimicrobials Reported:

Note: *E. coli* ATCC 25922 control organism. Acceptable ranges for antimicrobials, please refer to CLSI document M100-S18, Table 3, pg 142-143.

Antimicrobials	S	I	R
Ampicillin	≤ 8	16	≥ 32
TMP\SMX	≤ 2/38	-	≥ 4/76
Ciprofloxacin	≤1	2	≥ 4
Ceftriaxone	≤ 8	16-32	≥ 64
Chloramphenicol	≤ 8	16	≥ 32
Nalidixic Acid*	≤ 16	-	≥ 32

\*The following warning label is put if the extraintestinal *Salmonella* strain is resistant to Nalidixic acid.

This strain is resistant to Nalidixic acid. Fluoroquinolone-susceptible strains (i.e. Ciprofloxacin) of *Salmonella* that test resistant to Nalidixic Acid may be associated with clinical failure or delayed response in fluoroquinolone-treated patients with extraintestinal salmonellosis.



### 3.3 Pathogenic *E. coli*

#### Retail Meat

##### **Detection Assay**

All meat samples are screened for the presence of verocytotoxins using the verocell assay. This method is based on the MFLP-89 VTEC detection in the *Compendium of Analytical Methods, Health Protection Branch, Methods of Microbiological Analysis of Food, Government of Canada*. Fifty grams of meat sample is processed in 450 ml buffered peptone water (BPW) incubated at 35°C for 24 hrs and 1 mL is transferred into 9 ml of MacConkey broth. After incubation 100µl is transferred into brain heart infusion (BHI) broth and further incubated at 35°C for 6 hrs. The verocell assay is prepared and incubated at 35°C for 48 hrs. Microtitre plates are examined microscopically for cytotoxic activity. Confirmation of positive isolates is performed using biochemical methods.

##### ***E. coli* Coliform: Detection and Counts**

All meat samples are tested using the 3M Petrifilm™ *E. coli* / Coliform count plate. Meat samples are processed in 450 ml BPW; one millilitre is transferred onto Petrifilm™ and incubated at 35°C for 48 hrs. Bacterial colonies are counted and CFU/ml is calculated.

#### Agriculture

##### **Culture Detection**

All fecal samples are tested for the presence of *E. coli* 0157 using the Dynabead™ Anti-*E. coli* 0157 test. This method is based on the MFLP-90 and MFLP-80 Identification of *E. coli* 0157 by Dynabead™ anti-*E. coli* 0157 method of the *Compendium of Analytical Methods, Health Protection Branch, Methods of Microbiological Analysis of Food, Government of Canada*. Ten grams of fecal sample is pre-enriched in 90 ml BPW, homogenized and 1 ml is transferred to a 9 ml Brilliant Green Bile broth selective enrichment. The sample is incubated at 35°C for 6 hrs on a shaker. After incubating, 0.1 ml of sample is added to Dynabeads and further incubated at 25°C for 6 hrs on a shaker. The enrichment mixture is prepared for immunomagnetic separation (IMS) and plated onto three selective media (Rainbow agar, Cefixime Tellurite Sorbitol MacConkey (CT-SMAC) agar and CHROMagar O157) (since 2009, only CHROMagar selective medium has been used for plating). Samples are incubated at 35°C for 24 hrs and examined. Suspect colonies are further tested for a Tryptic Soy Blood agar slant using MacConkey broth and EC broth +MUG. Positive isolates are confirmed with slide agglutination using 0157 antisera.

##### **PCR Testing for Confirmation of :H7**

Presumptive *E. coli* 0157:H7 isolates are sent to The Escherichia coli Laboratory, (OIE reference laboratory), Faculté de médecine vétérinaire of University of Montreal for :H7 testing using polymerase chain reaction (PCR).

## **Untreated Surface Water**

### **Culture Detection**

All water samples are tested for the presence of *E. coli* O157 using Health Canada, Health Products and Food Branch using methods that comply with MFLP-80 "Isolation of *E. coli* O157:H7 or NM in Foods", supplements 1 and 2 to MFLP-80 and MFLP-90 "Identification of *E. coli* O157 by Dynabeads anti-*E. coli* O157" (LSD- MID 125 " Isolation and Identification of *E. coli* O157:H7 or NM in foods"). A 1L water sample is filtered using 0.45 µm filter membrane. Filter samples are enriched in 50 ml of non-selective TSB broth, incubated at 35°C for 4-6h. Novobiocin is added to a final concentration of 20 µg/ml and the enriched samples are further incubated at 42°C for 18-24h. After incubation, the samples are processed using Dynal Dynabeads™ Immunomagnetic separation (IMS) method (anti O157) and plated onto a selective media Cefixime Tellurite Sorbitol MacConkey (CT-SMAC) agar. Samples are incubated at 35°C, for 18-24h and examined. Suspect colonies are further tested using the *E. coli* O157 MUG test (using LST broth with methylumbelliferyl β glucouronide), cellobiose fermentation test, agglutination with *E. coli* O157 antisera, and API 20E test strip. Confirmation is conducted using the Assurance GDS for *E. coli* O157:H7 Gene Detection System (Health Canada method MFLP-16).

### **Subtyping (Retail, Agriculture, Water)**

#### **PFGE Testing**

Presumptive *E. coli* O157 isolates are sent to LFZ-Guelph for H7 typing and PFGE analysis. PFGE is performed at the Laboratory for Foodborne Zoonoses, Guelph (*E. coli* Lab). LFZ-protocols for *E. coli* serovar typing protocol and the *E. coli* PFGE protocols that comply with standardized protocols developed by PulseNet USA and in use by PulseNet Canada. For *E. coli*, only one restriction enzyme, *Xba*I, is used to obtain the PFGE pattern. The resulting tiff files are uploaded to the national database located at NML in Winnipeg to obtain the national designation for the two separate patterns.

#### **Antimicrobial Susceptibility Testing (2005-2007)**

Antimicrobial susceptibility testing is performed by the CIPARS AMR Laboratory, St-Hyacinthe using the Sensititre® Automated Microbiology System (Trek Diagnostic Systems Ltd.). *E. coli* isolates are prepared by streaking the cultures onto Mueller Hinton Agar for isolated colonies, and then for growth, and incubated for 18 to 24 hrs at 35°C. Each culture is prepared in Sensititre® demineralized water to a 0.5 McFarland standard and then inoculated into Sensititre® Mueller Hinton Broth with TES. The NARMS Gram negative plate with the CMV1AGNF panel is inoculated with the broth suspension and is incubated for 18 hrs at 35°C. Each plate is then read by the Sensititre® AutoReader to determine the minimum inhibitory concentration. The following are the antimicrobials tested: amikacin, ampicillin, amoxicillin/clavulanic acid, cefoxitin, ceftiofur, ceftriaxone, chloramphenicol, ciprofloxacin, gentamicin, kanamycin, nalidixic acid, streptomycin, sulfisoxazole, tetracycline, trimethoprim/sulfamethoxazole. CIPARS breakpoints are used to determine resistance profiles and can be found at:

[http://www.phac-aspc.gc.ca/ciparspicra/analys\\_e.html](http://www.phac-aspc.gc.ca/ciparspicra/analys_e.html).

## **Human**

### **LAB A**

Specimens are collected and mixed with Enteric Plus transport media before arriving at the laboratory. Once at the laboratory, a sterile swab is used to obtain a homogenized sample, which is streaked for isolated colonies onto Chromogenic O157 agar. Samples are incubated at 35°C for 18 to 24 hrs. Plates are examined for typical *E. coli* O157 colony morphology on both agar plates. One suspected colony is subcultured to a sheep blood agar if needed. Presumptive colonies are tested using the O157 Antisera and Microscan biochemical panel of tests. Confirmation of H7 is completed by TRPHL.

### **LAB B**

Specimens are collected and mixed with Cary-Blair transport media before arriving at the laboratory. Once at the laboratory, a sterile swab is used to obtain a homogenized sample, which is streaked for isolated colonies onto Sorbitol MacConkey (SMAC) agar. Samples are incubated at 35°C for 24 hrs. Plates are examined for typical *E. coli* O157 colony morphology and one suspected colony is subcultured onto a blood agar plate if needed. Presumptive colonies are tested using the O157 Antisera, Latex agglutination and the Vitek 2 ID system. Isolates are forwarded to TRPHL.

### **LAB C**

Specimens are collected and mixed with Cary-Blair transport media before arriving at the laboratory. Once at the laboratory, a sterile swab is used to obtain a homogenized sample which is streaked for isolated colonies onto Sorbitol MacConkey (SMAC) agar and Chromogenic agar. Samples are incubated at 36°C for 18 to 24 hrs. Plates are examined for typical *E. coli* O157 colony morphology on both agar plates. One suspected colony is picked from each plate and subcultured to a blood agar if needed. Presumptive colonies are tested using the Microgen O157 Antisera agglutination test and the Phoenix automated ID system.

### **LAB D**

Specimens are collected in a container with Cary-Blair transport media and mixed by the patient before arriving in the laboratory. Once at the laboratory, a sterile swab is used to obtain a homogenized sample, which is streaked for isolated colonies onto Sorbitol MacConkey (SMAC) agar. Samples are incubated at 37°C for 18 to 24 hours. Plates are examined for typical *E. coli* non-sorbitol fermenting colony morphology. One suspected colony is subcultured to a fresh SMAC agar plate or blood agar if needed. Presumptive colonies are tested using the O157 Antisera agglutination test and the Vitek automated ID system. Isolates are forwarded to TRPHL for 'H' agglutination.

## **Ontario Agency for Health Protection and Promotion, Public Health Laboratory – London and Toronto locations**

Stool specimens are collected and mixed with Cary-Blair transport medium at the time of collection. Once received in the laboratory, a sterile swab is used to obtain a representative portion of the sample which is streaked for isolated colonies onto a Sorbitol MacConkey Agar (SMAC) plate. In cases where *E.coli* O157:H7 is specifically suspected and requested a Cefixime-Tellurite Sorbitol MacConkey Agar (CT-SMAC) and MacConkey Broth are also inoculated. Inoculated media are incubated at 35°C for 18 -

24 hours. After incubation, a loopful of the MacConkey Broth is streaked for isolated colonies onto Sorbitol MacConkey Agar (SMAC) and CT-SMAC Agar plates. The plates are incubated at 35°C for 18-24 hours. Plates are examined for typical *E. coli* O157 non-sorbitol fermenting colony morphology and screened using TSI, ONPG-PA-M with indole strip and Lysine decarboxylase. Testing performed on presumptive colonies include the API 20E test and *E. coli* O157 antisera slide agglutination test from a Blood Agar plate. Further confirmatory testing performed at the Public Health Laboratory - Toronto include using a freshly sub-cultured Blood Agar plate, SMAC plate and MacConkey plate for testing isolates with a series of tube agglutinations for both the somatic (O157) and flagellar (H7) antigens as well as a panel of biochemical tests.

### **Subtyping (Human)**

Once confirmed positive, one isolate from each positive specimen is tagged as a C-EnterNet sample and sent to the Public Health Laboratory - Toronto for H7 typing and PFGE.

### **PFGE testing**

The PFGE method used at the Public Health Laboratory – Toronto is based on the national standardized PulseNet Canada protocol.

The one-day (24-48h) Standardized Laboratory Protocol for the Molecular Subtyping of Enteric Pathogens by Pulsed Field Gel Electrophoresis (PFGE) and Standardized Molecular Subtyping of Foodborne Bacterial Pathogens by Pulsed Field Gel Electrophoresis (Center for Disease Control and Prevention Manual –PNL 05) protocols are followed.

### **Antimicrobial Susceptibility Testing**

**Not done**

## 3.4 *Yersinia*

### Retail Meat

#### **Culture Detection**

Screening for *Yersinia enterocolitica* is performed on pork samples.

#### **Method # 1 (utilized in 2005-2008)**

Fifty grams of pork meat is processed in 450 ml of Luria Bertani-Bile Salt Irgasan (LB-BSI) broth, incubated at 12°C for 24 hrs and inoculated with 5ml of Irgasan solution. Samples are incubated at the same temperature for 48 hrs longer. Following incubation, LB-BSI is streaked onto selective agar plates and incubated at 30°C for 20 hrs. Suspect colonies are further screened and confirmed using biochemical methods.

#### **Method #2 (implemented in 2009)**

Fifty grams of pork meat is processed in 450 ml of Luria Bertani-Bile Salt Irgasan (LB-BSI) broth, incubated at 12°C for 24 hrs and inoculated with 5ml of Irgasan solution. Samples are incubated at the same temperature for 48 hrs longer. *Following incubation, 0.5 mL of LB-BSI broth is added to 4.5 mL of alkali solution and vortexed. The alkali-treated LB-BSI is streaked onto CIN agar plates and incubated at 30°C for 20 hrs.* Suspect colonies are further screened and confirmed using biochemical methods.

### Agriculture

#### **Culture Detection**

Screening for *Yersinia enterocolitica* is performed on swine manure samples. Twenty five grams of swine fecal sample is processed in 100 ml of BPW broth, homogenized and 5 ml is transferred to 100 ml of Irgasan Tircacillin Broth (ITCB). The sample is incubated at 24°C for 72 hrs. Following incubation, ITCB is streaked onto Cufsulodin-Irgasan Novobiocin (CIN) agar and incubated at 28°C for 48 hrs. Suspected colonies are further tested using TSI, MacConkey and Urea agar. Confirmation for *Yersinia enterocolitica* is performed using the API 20E test. One presumptive isolate is sent to TRPHL for serotyping and PFGE and to CIPARS AMR Laboratory, Saint-Hyacinthe Unit for antimicrobial susceptibility testing.

### Water

#### **Culture Detection**

#### **Method effective from the start of testing to Sept 9, 2008:**

A 1L water sample is filtered using 0.45µm filter membrane. Filter samples are enriched in *Yersinia* Trypticase Soy Broth (YTSB) and incubated at 12°C for 24±2h. Irgasan solution is then added and samples are incubated at the same temperature for 48h longer. From the enrichment broth one Cefsulodin-Irgasan-Novobiocin (CIN) agar plate is streaked and 0.5ml of broth is added to KOH/NaCl solution, followed by streaking to a second CIN plate. CIN plates are incubated at 30°C for 16-18h.

Suspect isolates are further tested using oxidase tests, MacConkey, Triple Sugar Iron (TSI) and Christensen's urea agar. Optional confirmation is performed using API 20E test when warranted.

**Method effective from Sept 10, 2008 – Jan 7, 2009:**

A 1L water sample is filtered using 0.45µm filter membrane. Filter samples are enriched in Luria-Bertani-Bile Salts Irgasan (LB-BSI) Broth with CN Supplement and incubated at 12°C for 24±2h. Irgasan solution is then added and samples are incubated at the same temperature for 48h longer. From the enrichment broth one Cefsulodin-Irgasan-Novobiocin (CIN) agar plate is streaked and 0.5ml of broth is added to KOH/NaCl solution, followed by streaking to a second CIN plate. CIN plates are incubated at 30°C for 16-18h.

Suspect isolates are further tested using oxidase tests, MacConkey, Triple Sugar Iron (TSI) and Christensen's urea agar. Optional confirmation is performed using API 20E test when warranted.

**Method effective from Jan 8, 2009 – current:**

A 1L water sample is filtered using 0.45µm filter membrane. Filter samples are enriched in Yersinia Trypticase Soy Broth (YTTSB) and incubated at 12°C for 24±2h. Irgasan solution is then added and samples are incubated at the same temperature for 48h longer. From the enrichment broth one Cefsulodin-Irgasan-Novobiocin (CIN) agar plate is streaked and 0.5ml of broth is added to freshly prepared (7 day shelf life) KOH/NaCl solution, followed by streaking to a second CIN plate. CIN plates are incubated at 30°C for 16-18h in a moisture chamber.

Suspect isolates are further tested using oxidase tests, MacConkey, Simmon's citrate, Triple Sugar Iron (TSI), Christensen's urea agar, and Motility at 25°C and 37°C. Optional confirmation is performed using API 20E test when warranted.

Susceptibility testing is performed by LFZ-SHU (LFZ-SHU Isolation protocol). Serotyping analysis is conducted by TRPHL.

**Subtyping (Retail, Agriculture, Water)**

**Serotyping, Biotyping**

Presumptive *Yersinia enterocolitica* isolates from the retail, agriculture and water samples are sent to TRPHL for speciation, serotyping and biotyping. The serotyping and biotyping methods for all *Yersinia* isolates are based on the methods of Wauters (1981), Wauters (1991), Bottone (1997), Bockemuhl (2003) and Aleksic (1984).

**Antimicrobial Susceptibility Testing (2005-2007)**

Antimicrobial susceptibility testing is performed by the CIPARS AMR Laboratory, St-Hyacinthe Unit using the Sensititre® Automated Microbiology System (Trek Diagnostic Systems Ltd.). *Yersinia* isolates are prepared by streaking the cultures onto Mueller Hinton Agar for isolated colonies, and then for growth, and incubated for 18 to 24 hrs at 35°C. Each culture is prepared in Sensititre® demineralised water to a 0.5 McFarland standard and then inoculated into Sensititre® Mueller Hinton Broth with TES. The NARMS Gram negative plate with the CMV1AGNF panel is inoculated with the broth suspension and is incubated for 18 hrs at 35°C. Each plate is then read by the Sensititre® AutoReader to determine the minimum inhibitory concentration. The following are the antimicrobials tested: amikacin, ampicillin, amoxicillin/clavulanic acid, cefoxitin,

ceftiofur, ceftriaxone, chloramphenicol, ciprofloxacin, gentamicin, kanamycin, nalidixic acid, streptomycin, sulfisoxazole, tetracycline, trimethoprim/sulfamethoxazole. CIPARS breakpoints for *E. coli* are used to determine resistance profiles and can be found at: [http://www.phac-aspc.gc.ca/cipars-picra/analys\\_e.html](http://www.phac-aspc.gc.ca/cipars-picra/analys_e.html).

### **Enumeration - Most Probable Number (MPN)**

The Most Probable Number method is used for retail pork samples positive for *Yersinia enterocolitica* from the initial primary detection method. The MPN method estimates the number of *Yersinia* species per millilitre or gram of sample using three tube dilutions. Fifty millilitres of the original processed sample is stored for subsequent MPN testing and used to prepare serial dilutions. Tubes are incubated at 12°C for 24 hrs. After incubation, 100µl of an Irgasan solution is added to each tube and they are incubated at 12°C for 48 hrs. Then each tube is streaked onto selective agar plates and incubated at 30°C for 20 hrs. Suspect colonies are further screened and confirmed using biochemical methods. MPN is calculated using the MPN table in the FDA Bacteriological Analytical Manual (<http://www.cfsan.fda.gov/~ebam/bam-toc.html>).

## **Human**

### **LAB A**

Specimens are collected and mixed with Enteric Plus transport media before arriving at the laboratory. Once at the laboratory, a sterile swab is used to obtain a homogenized sample, which is streaked for isolated colonies onto SS agar and incubated at 35°C for 18 to 48 hrs. Presumptive *Yersinia* spp. identified as small NLF colonies is further tested using Microscan, TSI, OPM and motility testing for confirmation. All *Yersinia* species are sent to TRPHL for typing.

### **LAB B**

*Yersinia enterocolitica* testing is only performed routinely for children <12yrs of age. Specimens are collected and mixed with Cary-Blair transport media before arriving at the laboratory. Once at the laboratory, a sterile swab is used to obtain a homogenized sample which is streaked for isolated colonies onto *Yersinia* selective agar (CIN) and incubated at 30°C for 24 hrs. Presumptive *Yersinia* spp. are identified as small, dark pink “bull’s eye” colonies, and confirmation is done with Triple Sugar Iron (TSI) and the Vitek 2 ID system. All *Yersinia* species are sent to TRPHL for typing.

### **LAB C**

*Yersinia enterocolitica* testing is only performed routinely for children <12yrs of age. Specimens are collected and mixed with Cary-Blair transport media before arriving at the laboratory. Once at the laboratory, a sterile swab is used to obtain a homogenized sample which is streaked for isolated colonies onto *Yersinia* selective agar (CIN) and incubated at room temperature for 48 hrs. Presumptive *Yersinia* spp. are identified as small, dark pink “bull’s eye” colonies, and confirmation is done with TSI, Urea, SIM and the Phoenix automated ID system.

## **LAB D**

Specimens are collected in a container with Cary-Blair transport media and mixed by the patient before arriving at the laboratory. Once at the laboratory, a sterile swab is used to obtain a homogenized sample, which is streaked for isolated colonies onto *Yersinia* selective agar (CIN) and incubated at 30°C for 24 hours. Presumptive *Yersinia* spp. are identified as small, dark pink “bull’s eye” colonies, and confirmation is done with Catalase, Salicin and the Vitek automated ID system. Isolates are forwarded to TRPHL for typing.

### **Ontario Agency for Health Protection and Promotion, Public Health Laboratory – London and Toronto locations**

Stool specimens are collected and mixed with Cary-Blair transport medium at the time of collection. Once received in the laboratory, a sterile swab is used to obtain a representative portion of the sample which is streaked for isolated colonies onto a Cefsulodin-Irgasan-Novobiocin Agar (CIN) plate. The CIN plate is incubated at 25-32°C for 18-24 hours. Presumptive *Yersinia* spp. are identified as small, dark pink “bull’s eye” colony and screened using TSI, ONPG-PA-M with indole strip, Lysine decarboxylase and an API 20E test. Further confirmatory testing performed at the Public Health Laboratory - Toronto include using a freshly sub-cultured Blood Agar plate, MacConkey plate and Congo-Red Magnesium Oxalate Agar (CRMOX) for performing serotyping as well as a panel of biochemical tests for biotyping.

### **Subtyping (Human)**

*Yersinia enterocolitica* isolates are tagged with a C-EnterNet label and submitted to Public Health Laboratory - Toronto for serotyping and biotyping.

### **Serotyping and Biotyping Method References**

The serotyping and biotyping methods of all *Yersinia* isolates are based on the following references:

Wauters G. 1981. Antigens of *Yersinia enterocolitica*, pp. 41-53. *In: Yersinia enterocolitica*. EJ Bottone (ed). CRC Press, Boca Raton, FL

Clinical Microbiology Review, Apr. 1997, p. 257-276.

Wauters G et al. 1991 Somatic and flagellar antigens of *Yersinia enterocolitica* and related species. *Contrib. Microbiol. Immunol.* 12: 239-243.

Bockemuhl J, Wong DJ. 2003 *Yersinia* In Murray PR *et al* (ed), Manual of Clinical Microbiology, 8<sup>th</sup> ed, p. 672, ASM Press, Washington, DC.

Aleksic S, Bockemuhl J. 1984. Proposed revision of the Wauters et al. antigenic scheme for serotyping of *Yersinia enterocolitica* *J. Clin. Microbiol.* 20 (1): 99-102.

### **Antimicrobial Susceptibility Testing**

Not done



## 3.5 *Listeria*

### **Retail Meat**

#### **Culture Detection**

Samples are processed in *Listeria* Enrichment Broth (LEB) and incubated at 30°C for 24 and 48 hrs. This method is based on the MFHPB-30 *Listeria monocytogenes* detection in the *Compendium of Analytical Methods, Health Protection Branch, Methods of Microbiological Analysis of Food, Government of Canada*. After enrichment incubation, 0.1ml of LEB is transferred to Modified Fraser Broth (MFB) and incubated at 35°C for 48 hrs. MFB inoculum is streaked onto selective agar plates, incubated at 30°C for 24 to 48 hrs. Suspect colonies are screened and confirmed using biochemical methods and microscopy.

### **Agriculture**

#### **Culture Detection**

Samples are processed by transferring one part fecal sample into nine parts *Listeria* Enrichment Broth (LEB) and incubated at 30°C for 24 to 48 hrs. This method is based on the MFHPB-30 *Listeria monocytogenes* detection in food method of the *Compendium of Analytical Methods, Health Protection Branch, Methods of Microbiological Analysis of Food, Government of Canada*. The complete method is performed for LEB broth after 24 hrs incubation and separately for 48 hrs incubated broth. After enrichment incubation, 0.1ml of LEB is transferred to Modified Fraser Broth (MFB) and incubated at 35°C for 48 hrs. MFB inoculums is streaked onto Oxford Agar (OXA Palcam agar and Rapid'L.mono agar plates (chromogenic agar) and incubated at 30°C for 24 to 48 hrs and suspect colonies are further tested on MHB agar. Biochemical tests are performed using the method MFHPB-30 list of tests as a guideline and include motility testing, mannitol, rhamnose, xylose sugar utilization and a Gram stain. Biochemical testing plates are incubated at 35°C for 48 hrs. Confirmation testing is done using slide agglutination with *Listeria* antiserum.

### **Subtyping (Retail, Agriculture)**

#### **Confirmation typing and PFGE**

*Listeria monocytogenes* isolates are further tested using the PFGE methodology outlined by Graves (2001).

#### **Enumeration- Most Probable Number (MPN)**

The Most Probable Number method is only used for retail samples positive for *Listeria monocytogenes*. This method estimates the number of *Listeria* species per ml or g of sample using three tube MPN dilutions. Fifty millilitres of the original processed sample is stored for MPN testing and used to prepare serial dilutions. Tubes are incubated at 30°C for 48 hrs. After incubation, 100µl from each tube is transferred to 10 ml of MFB and incubated at 35°C for 24 hrs. Then, MFB inoculums are streaked onto selective agar plates and incubated at 30°C for 24 to 48 hrs. Suspect colonies are screened and confirmed using biochemical methods and microscopy.

MPN is calculated using the MPN from the FDA Bacteriological Analytical Manual (<http://www.cfsan.fda.gov/~ebam/bam-toc.html>).

## **3.6 *Giardia* and *Cryptosporidium***

### **Agriculture**

During the first year of sampling each commodity is tested for the presence of *Giardia* and *Cryptosporidium*. Fecal floats are prepared for DNA extraction and microscopy. A Two-step nested PCR analysis is conducted for *Giardia* and *Cryptosporidium*. For *Giardia*, round one primers used are Gia2029 and Gia2150c, followed by RH4R and RH11F in round two of testing. For *Cryptosporidium*, the round one primers are 1325R and 1325R, followed by 826F and 826R for round two of testing.

### **Untreated Surface Water**

#### **Microscopic Detection**

Testing for *Giardia* and *Cryptosporidium* is performed on water samples once per month by Hyperion Research Ltd. A 25 litre river water sample is passed through a Filta-max filter (Envirochek) in the field, and the filter is then shipped to Hyperion Research Ltd. and processed for parasite enumeration using immunomagnetic separation followed by fluorescence microscopy (USEPA Method 1623 *Cryptosporidium* and *Giardia* in water by filtration/IMS/FA).

#### **Molecular Typing**

*Cryptosporidium* genotyping is done by removing the coverslips and mounting media from positive slides. Material is then removed from the entire well using four rinses with lysis buffer and scraping with a pipette tip. DNA is extracted by repeated freeze-thawing of oocysts and purified using the QIAGEN DNA Micro Kit. Nested PCR is done by targeting the small ribosomal subunit RNA (18S rRNA) gene with primary and secondary primers as described by Xiao *et al.* (1999). DNA sequencing of the secondary PCR product is used to confirm *Cryptosporidium* species/genotypes. Bidirectional sequence analysis is performed using an ABI Prism 3130xl Genetic Analyzer (Applied Biosystems).

### **Human**

#### **Lab D**

Stool concentrates are prepared for microscopic identification by placing a drop of the concentrate on a glass slide, diluting it with saline or formalin if necessary, and placing a cover slip over top of the drop. Stool received unfixed are stained with modified acid fast/Iron Haematoxylin. The entire slide is scanned on low power (10x) looking for ova and larvae. Cysts and trophozoites are scanned using a 50x oil immersion objective. A 100x oil immersion objective scan is used to identify smaller cysts and trophozoites i.e. *Enteromonas hominis*. All organisms are measured with a calibrated ocular micrometer.

**Ontario Agency for Health Protection and Promotion,  
Public Health Laboratory – Toronto**

SAF preserved fecal specimens are processed by the following method for ova and parasites including *Giardia* and *Cryptosporidium*.

Routine processing of SAF-preserved fecal specimen involves the following steps: Firstly, an initial processing is carried out in which a portion of the specimen is filtered through gauze and washed twice with physiological saline to produce a sediment. From this sediment, a small amount is retrieved and mixed with gelatin adhesive to make a smear for iron-haematoxylin staining. Then, the remainder of the sediment is mixed with formalin, then di-ethyl ether and centrifuged to produce a concentrate after the plug, the aqueous and organic material are decanted. The stained smear and concentrate are used for the recovery of most helminth eggs/larvae and protozoan cysts/trophozoites including *Giardia*. In addition to these two items, directly from the specimen two air-dried smears are prepared. One is stained with the fluorescent stains auramine-rhodamine for the detection of *Cryptosporidium* and other coccidians. The second smear is stained with modified trichrome stain for microsporidia spores. Additional testing such as direct wet mounts and other concentration procedures are also performed depending on the parasite suspected.

## 3.7 Viruses

### Agriculture

During the first year of sampling each commodity is tested for the presence of rotavirus and norovirus. Fecal samples are filtered and prepared for PCR analysis using the one-step RT-PCR reactions. Presumptive positives are gel purified and sequenced on the tested strands.

Fecal samples are suspended at 5% w/v in 0.9% NaCl, vortexed briefly and clarified through a combination glass fiber/PVDF 0.22  $\mu$ M filter. RNA is extracted from a 140  $\mu$ l sample of the resulting filtrate using the QIAamp viral RNA extraction kit according to the manufacturer's recommendations. Total RNA is extracted from 25 g meat samples using Tri reagent and Dynabeads coated with oligo-dT as described by Kinglsey and Richards (2001).

The RNA was used as a template for one-step RT-PCR reactions using Monroe region B primers (Anderson et al., 2003) or Rota1 and End9 primers (Gouvea et al., 1990; Jean et al., 2002) and the Qiagen One Step RT-PCR kit. 213 bp (norovirus) and 268 bp (rotavirus) amplicons are considered to be presumptively positive. They are gel purified using a QIAquick gel extraction kit and sequenced on both strands by DNA Landmarks.

### 3.8 Sources Cited

- Aleksic S, Bockemuhl J. 1984. Proposed revision of the Wauters et al. antigenic scheme for serotyping of *Yersinia enterocolitica*. *J Clin Microbiol* 20 (1): 99-102.
- Anderson ES, Williams REO. 1956. Bacteriophage typing of enteric pathogens and staphylococci and its use in epidemiology. *J Clin Pathol* 9:94-114.
- Anderson ES. 1964. The phage typing of *Salmonella* other than *S. typhi*. In: Van Oye E, ed. The world problem of salmonellosis. The Hague, The Netherlands: Dr W. Junk Publishers: 89-100.
- Anderson ES, Ward LR, de Saxe MJ, de Sa JDH. 1977. Bacteriophage-typing designations of *Salmonella* Typhimurium. *J Hyg* 78:297-300.
- Anderson AD et al 2003. A Waterborne Outbreak of Norwalk-Like Virus among Snowmobilers—Wyoming, 2001. *J Infect Dis* 187: 303-6.
- Bhaduri S, Wesley IV, and Bush EJ. 2005. Prevalence of Pathogenic *Yersinia enterocolitica* Strains in Pigs in the United States. *Appl Environ Microbiol* 71:7117-7121.
- Bockemuhl J, Wong DJ. 2003. *Yersinia*. In: Murray PR et al ed., Manual of Clinical Microbiology, 8<sup>th</sup> ed, p. 672, ASM Press, Washington, DC.
- Bottone, EJ. 1997. *Yersinia enterocolitica*: the charisma continues. *Clin Microbiol Rev* 10:257-276.
- Callow BR. 1959. A new phage typing scheme for *Salmonella typhimurium*. *J Hyg*. 57:346-59.
- Clinical and Laboratory Standards Institute (CLSI). *Performance Standards for Antimicrobial Susceptibility Testing; Eighteenth Informational Supplement*. CLSI document M100-S18. Wayne, PA: Clinical and Laboratory Standards Institute: 2008.
- Demczuk W, Soule G, Clark C, Ackermann, Hans-W, Easy R, Kahkhria R, Rodgers F, Ahmed R. 2003. Phage-based typing scheme for *Salmonella enterica* serovar Heidelberg, a causative agent of food poisonings in Canada. *J Clin Microbiol* 41: 4279-4284.
- Ewing WH. Edwards and Ewing's Identification of *Enterobacteriaceae*, 4<sup>th</sup> ed, pp. 181-304, 1986, Elsevier Science Publ., New York.
- Farmer III JJ. 2003 *Enterobacteriaceae*: introduction and Identification, p. 636-653. In PR Murray, EJ Baron, MA Pfaller, JH Jorgensen, and R H Tenover ed., Manual of clinical microbiology, 8<sup>th</sup> ed. American Society for Microbiology, Washington, D.C.
- Gouvea V, Glass RI, Woods P, Taniguchi K, Clark HF, Forrester B, and Fang ZY. 1990. Polymerase chain reaction amplification and typing of rotavirus nucleic acid from stool specimens. *J Clin Microbiol* 28: 276-82.
- Graves LM, Swaminathan B. 2001. PulseNet standardized protocol for subtyping *Listeria monocytogenes* by macrorestriction and pulsed-field gel electrophoresis. *Int J Food Microbiol* 65:55-62.

Jean J,Blais B, Darveau A,Fliss I. 2002. Simultaneous detection and identification of hepatitis A virus and rotavirus by multiplex nucleic acid sequence-based amplification (NASBA) and microtiter plate hybridization system. *J Virol Methods* 105:123-32.

Iijima Y, Asako NT, Aihara M, Hayashi K. 2004. Improvement in the detection rate of diarrhoeagenic bacteria in human stool specimens by a rapid real-time PCR assay. *J Med Microbiol* 53:617-622.

Kingsley DH,Richards GP. 2001. Rapid and Efficient Extraction Method for Reverse Transcription-PCR Detection of Hepatitis A and Norwalk-Like Viruses in Shellfish. *Appl Environ Microbiol* 67: 4152-7.

Le Minor L, Popoff MY. 2001. Antigenic formulas of the *Salmonella* serovars. 8th ed., WHO Collaborating Centre for Reference and Research on *Salmonella*, Paris.

Lübeck PS, Wolffs P, On SLW, Anhrens P, Rådström P,Hoorfar J. 2003. Toward an international standard for PCR-based detection of food-borne thermotolerant *Campylobacters*: assay development and analytical validation. *Appl Environ Microbiol* 69:5664-5669.

Popoff MY. 2001. Antigenic Formulas of the *Salmonella* Serovars, 8<sup>th</sup> ed., 2001. WHO Collaborating Center for Reference and Research in *Salmonella*, Institut Pasteur, Paris, France.

Ram S, Shanker R. 2005. Computing Taqman probes for multiplex PCR detection of *E. coli* O157 serotypes in water. *Silico Biology* 5:0045.

Rodríguez-Lázaro D, Hernández M, Esteve T, Hoorfar J,Pla M. 2003. A rapid and direct real time PCR-based method for identification of *Salmonella* spp. *J Microbiol Meth* 54:381-390.

Sharma VK,Dean-Nystrom EA. 2003. Detection of enterohemorrhagic *Escherichia coli* O157:H7 by using a multiplex real-time PCR assay for genes encoding intimin and Shiga toxins. *Vet Microbiol* 93:247-260.

Shipp CR Rowe B. 1980. A mechanised microtechnique for salmonella serotyping. *J Clin Path* 33: 595-597.

Ward LR, de Sa JDH, Rowe B. 1987. A phage-typing scheme for *Salmonella enteritidis*.*Epidemiol Infect* 99:291-294.


Wauters G. 1981. Antigens of *Yersinia enterocolitica*, pp. 41-53. *In: Yersinia enterocolitica*. EJ Bottone (ed). CRC Press, Boca Raton, FL

Wauters G et al. 1991, Somatic and flagellar antigens of *Yersinia enterocolitica* and related species. *Contrib Microbiol Immunol* 12:239-243

Xiao, L, Morgan UM, Limor J, Escalante A, Arrowood M,Shulaw W, Thompson RC, Fayer R, Lal AA. 1999. Genetic Diversity within *Cryptosporidium parvum* and Related *Cryptosporidium* Species. *Appl Environ Microbiol* 65:3386-3391.




# C-EnterNet Retail Sample Collection Overview





# C-EnterNet Agriculture Sample Collection Overview








# C-EnterNet Watershed Data Collection Overview

C-EnterNet SOP Manual -Water





# C-EnterNet Human Sample Collection Overview

